



Updated July 2019

Patient Name, Today's Date, NEW Patient, CURRENT Patient, DOB, Height, Weight, Male, Female, Preferred Language, Best Phone, Email, Street Address, Apt#, City, State, Zip, Ship to Patient at: Home, Physician Office, Work Address, Allergies, Current Medications including OTC's (please fax a complete list)

Please Fax Insurance Card(s) both sides

Insured's Name, Relation to Patient, Primary Insurance, ID#, Group #, Secondary Insurance, ID#, Group #

Ordering Prescriber

Office Contact, Street Address, Suite #, City, State, Zip, Tel, Fax, Email, License#, NPI#

- ICD-10 Code C90.0 Multiple Myeloma, C91.10 Lymphoid Leukemia, D80.0 Hereditary Hypogammaglobulinemia, D80.1 Nonfamilial Hypogammaglobulinemia, D80.2 Selective deficiency of IgA, D80.3 Selective deficiency of IgG Subclasses, D80.5 Immunodeficiency with Increased IgM, D81.1 SCID with Low T- and B- Cell Numbers, D81.2 SCID with Low or Normal B-Cell Numbers, D81.9 Combined Immunodeficiency, Unspecified, D83.1 CVID w/ Predominant Immunoregulatory T-Cell Disorders, D83.8 Other Common Var. Immunodeficiencies, D83.9 Common Var. Immunodeficiency, Unspecified, E13.40 Other specified diabetes mellitus w/ diabetic neuropathy, unspecified, G25.82 Stiff Person Syndrome, G61.0 Guillain-Barre Syndrome (GBS), G62.9 Other Peripheral Neuropathy, G70.0 Myasthenia Gravis (MG), G70.80 Lambert-Eaton Syndrome, unspecified, L10.9 Pemphigus, M32.9 Systemic lupus erythematosus (SLE), M33.90 Dermatopolymyositis & Organ Involvement Unspecified, Q81.9 Epidermolysis Bullosa, Z94.81 BMT, C90.1 Plasma Cell Leukemia, D69.6 Thrombocytopenia, D80.1 Nonfamilial Hypogammaglobulinemia, D80.3 Selective deficiency of IgG Subclasses, D81.1 SCID with Low T- and B- Cell Numbers, D81.89 Other combined Immunodeficiencies, D83.1 CVID w/ Predominant Immunoregulatory T-Cell Disorders, D83.9 Common Var. Immunodeficiency, Unspecified, D84.9 CVID, G35 Multiple Sclerosis (MS), G61.81 Chronic Inflammatory Demyelinating Polyneuropathy (CIDP), G63 Polyneuropathy in diseases classified elsewhere, G70.01 Myasthenia Gravis with (Acute) Exacerbation, L12.0 Pemphigoid, M30.3 Kawasaki's syndrome, M33.20 Polymyositis, Organ Involvement Unspecified, M36.0 Dermatomyositis, Z41.8 Prophylactic Immunotherapy, Other:

PRESCRIPTION

SUPPLIES FOR INFUSION (If Necessary)

- NaCl 0.9% / D5W for flush: flush Line/Port with (3 - 5 ml for PIV and 5-10 ml for Central Line/Port) per nursing agency protocol (NaCl 0.9% / D5W will be used based on IVIG compatibility)
Heparin for flush (100 Units / ml) (if RN keeps PIV or if needed for Central Line), flush with 3-5 ml per nursing agency protocol
Sterile water for reconstitution of powder to make the requested concentration (for Carimune NF)
Other:

PRE-MEDICATIONS: To be administered 30 min prior to IVIG Infusion: (QTY: per infusion): Acetaminophen 650 mg PO

- Diphenhydramine 25 mg-50 mg Other:

IVIG (IMMUNOGLOBULIN) ORDER: (IVIG brand will be chosen if not specified)

INTRAVENOUS IMMUNOGLOBULIN Dose 0.4 gm/kg, 1gm/kg, 2gm/kg, gm

Infuse: IV daily for day(s); repeat every week(s) for cycles
Other:

Refills:
Refills:

SUBCUTANEOUS IMMUNOGLOBULIN Infuse: gm OR ml using sites time(s) per week for

Hydration order: ml NS IV to be infused prior/concurrently with IVIG

Refills:

ACCESS

- Peripheral
Midline, central (non-port), PICC
Implanted Port
Tunneled
Groshong PICC, Midline

NS HEPARIN

- 1-3ml before/after use
NS 5-10 mls before/after use;
5-10mls before/after use;
5-10mls before/after use;
5-10mls before/after use;

100 u/ml (If applicable, flush IV access device per Pharmacy protocol)

- 10u/ml 1-2mls after last NS flush
10mls after blood draw 10 u/ml 3-5mls after last NS flush; 5mls after blood draw
20mls after blood draw 100 u/ml 5mls after last NS flush; 5mls after blood draw
20mls after blood draw 10 u/ml 3- mls after last NS flush. 5mls after blood draw
10mls after blood draw NO Heparin needed

IN THE EVENT OF ANAPHYLAXIS:

- Stop Infusion and call MD & 911
Diphenhydramine 25 - 50 mg IVP every 4 hours prn (Not to exceed 25 mg/min) QTY: 3 (50 mg)
Epinephrine (1:1000) 0.4 mg SQ prn anaphylaxis, may repeat every 20 minutes x 2 QTY: 3 amp
Other:

ENROLL IN NURSE TRAINING / MANUFACTURER PROGRAM

Prescriber's Signature (signature required. NO STAMPS) Date

My signature certifies that the person named on this form is my patient and that the information provided on this enrollment form is complete and accurate to the best of my knowledge. I certify this therapy to be medically necessary.

My signature authorizes The Pharmacy and its representatives to act as the agent to execute the insurance prior authorization process, assist the above named patient enroll into patient support programs, and appeal on behalf of prescriber and patient in the event of a prior authorization denial.

IMPORTANT NOTICE: This fax is intended to be delivered only to the named addressee. It contains material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If you are not the named addressee, you should not disseminate, distribute, or copy this fax. Please notify the sender immediately if you have received this document in error and then destroy this document immediately.

PLEASE NOTE: The Pharmacy can only accept original prescription drug orders from patients, faxed prescriptions can be accepted only from the prescribing practitioners.